# TURTUR CAROLINENSIS.

## *The Turtle of Carolina.*

THIS is ſomewhat leſs than a Dove-houſe Pigeon: the Eyes black, compaſſed with a blew Skin: the Bill black: the upper part of the Head, Neck, Back, and upper part of the Wings brown: the ſmall feathers of the Wing, next the Back, have large black ſpots: the lower part of the Wing and Quill-feathers of a lead colour, three or four of the longeſt being almoſt black: The Breaſt and Belly of a pale Carnation colour. On each ſide the Neck, the breadth of a Man's Thumb, are two ſpots of the colour of burniſhed Gold, with a tincture of crimſon and green; between which and its Eye is a black ſpot. The Wings are long, the Tail much longer, reaching almoſt five Inches beyond them, and hath fourteen feathers, the two middle longeſt, and of equal length, and all brown; the reſt are gradually ſhorter, having their upper part lead colour, the middle black, and the end white. The Legs and Feet are red. They breed in *Carolina*, and abide there always. They feed much on the Berries of Poke, i. e. *Blitum Virginianum*, which are Poiſon. They likewiſe feed on the Seeds of this Plant; and they are accounted good Meat.

## Tourterelle de la Caroline.

*ELLE eſt un peu moins groſſe qu'un Pigeon domeſtique: ſes yeux ſont noirs, entourés d'une peau bleuë; ſon bec eſt noir; le deſſus de ſa tête, de ſon cou, de ſon dos, & la partie de ſes ailes plus proche des épaules, ſont bruns; les petites plumes des ailes les plus proches du dos ont de grandes taches noires; les autres plumes des ailes grandes & petites ſont couleur de Plomb; les trois ou quatre plus grandes ſont preſque noires; ſa poitrine & ſon ventre ſont d'une couleur de Roſes pâles; de chaque côté de ſon cou il y a une tache de largeur du pouce couleur d'or bruni mêlé de Cramoiſi & de Verd. Entre cette tache & l'aîl il y en a une autre noire; ſes ailes ſont longues; ſa queuë les paſſe de prés de cinq pouces: elle eſt compoſée de quatorze plumes; les deux du milieu ſont égales, plus longues que les autres & toutes brunes; les autres ſont toûjours plus courtes à meſure qu'elles s'eloignent des plumes du milieu; elles ſont blanches à l'extremité, noires au milieu, & couleur de Plomb en haut. Ses piés & ſes jambes ſont rouges. Ces oiſeaux ſont leurs Petits à la* Caroline, *& y demeurent toujours. Ils ſe nourriſſent de Bayes de Blettes de la* Virginie *qui ſont venimeuſes. Ils ſe nourriſſent auſſi des ſemences de la Plante décrite ci-deſſous; ſes ſemences ne ſont pas vénimeuſes.*

*Anapodophyllon Canadenſe Morini.* Turnef. Ranunculi facie planta peregrina H. R. Par. Aconitifolia humilis, flore albo, unico campanulato Fructu Cynosbati. *Mentz. Tab.* 11. *Turnef. Inſt. p.* 239.

## THE MAY APPLE.

THIS Plant grows about a Foot and half high; the Flower conſiſting of ſeveral *Petala*, with many yellow *Stamina* ſourrunding the ſeed-veſſel, which is oval, *unicapſular*, and contains many roundiſh Seeds. The Leaves of the Plant reſemble the *Aconitum lycoctonum luteum C. B. Pin.* The Root is ſaid to be an excellent Emetic, and is uſed as ſuch in *Carolina*; which has given it there the Name of *Ipecacuana*, the ſtringy Roots of which it reſembles. It flowers in *March*, the Fruit is ripe in *May*; which has occaſioned it in *Virginia* to be called *May-Apple*.

## Pomme de May.

*CETTE Plante s'éleve juſqu'à la hauteur d'un pié & demi. Sa fleur eſt compoſée de pluſieurs feuilles & de pluſieurs Etamines jaunes qui entourent l'Ovaire: Il eſt ovale & n'a qu'une ſeule Coſſe remplie de ſemences preſque rondes. Les feuilles de cette Plante ſont aſſés ſemblables à celles de l'Aconit Lycoctone jaune. On dit que ſa racine eſt un excellent Emétique; & l'on s'en ſert à la Caroline pour faire vomir; ce qui lui a fait donner dans ce païs là le nom d'*Ipecacuana, *outre qu'elle reſſemble aux racines fibreuſes de cette derniére. Cette Plante fleurit au mois de Mars, ſon fruit eſt meur en* May, *c'eſt pourquoi à la* Virginie *on l'apelle* Pomme de May.